

## Femoral neck fractures in patients with stroke sequelae

\*Ciobanu Natalia<sup>1,2</sup>, Ciobanu Sergiu<sup>1,3</sup>

<sup>1</sup>Department of Neurology No 2, Nicolae Testemitsanu State University of Medicine and Pharmacy

<sup>2</sup>Epilepsy and Cerebrovascular Diseases Laboratory, Institute of Emergency Medicine

<sup>3</sup>Clinical Hospital of Traumatology and Orthopedics, Chisinau, the Republic of Moldova

\*Corresponding author: nataliaandronic@yahoo.com. Received February 06, 2017; accepted April 03, 2017

### References

- Sacco R, Benjamin H, Broderick J, Dyken M, Easton J, American Heart Association Prevention Conference IV. Prevention and Rehabilitation of Stroke. Risk factors. *Stroke*, 1997.
- Gavriliuc M, Groppa S, Moldovanu I. Protocol clinic național. Accidental Vascular Cerebral Ischemic, Chișinău 2008.
- International Osteoporosis Foundation. Facts and statistics: <https://www.iofbonehealth.org/facts-statistics>.
- Andersson A. G, Seiger A, Appelros P. Hip fractures in persons with stroke. *Stroke research and treatment*, 2013.
- Peszczynski M. The fractured hip in hemiplegic patients. *Geriatrics*. 1957; 12: 687–90.
- Jacobsen B, Wilsgaard T. et al. Walking after stroke: does it matter? Changes in bone mineral density within the first 12 months after stroke: a longitudinal study. *Osteoporos. Int.* 2000; 11: 381–7.
- Ramnemark A. et.al, Fractures after stroke. *Osteoporos. Int.* 1998; 8: 92–5.
- Forster A, Young J. Incidence and consequences of falls due to stroke: a systematic inquiry. *BMJ*. 1995; 311: 83–6.
- Poole Kenneth, Jonathan Reeve, Elizabeth A. Warburton. Falls, fractures, and osteoporosis after stroke. *Stroke* 33.5 (2002): 1432-6.
- Ramnemark A, Nilsson M, Borsen B. et al. Stroke, a major and increasing risk factor for femoral neck fracture. *Stroke*. 2000; 31: 1572–7.
- Kanis J, Oden A. Acute and long-term increase in fracture risk after hospitalization for stroke. *Stroke*. 2001; 32:702–6.
- Dennis M, McDowell M. et al. Fractures after stroke: frequency, types, and associations / *Stroke*. 2002; 33: 728–34.
- Nevitt M, Cummings S. Type of fall and risk of hip and wrist fractures: the study of osteoporotic fractures. *J. Am. Geriatr. Soc.* 1993; 41: 1226–34.
- Chiu K. et al. A prospective study on hip fractures in patients with previous cerebrovascular accidents. *Injury*. 1992; 23: 297–99.
- Jorgensen L, Engstad T, Jacobsen B. Bone mineral density in acute stroke patients: low bone mineral density may predict first stroke in women. *Stroke*. 2001; 32: 47–51.
- Ramnemark A, Nyberg L. et al. Progressive hemosteoporosis on the paretic side and increased bone mineral density in the nonparetic arm the first year after severe stroke/ *Osteoporos. Int.* 1999; 9: 269–75.
- Price R. el. al, Forearm bone loss in hemiplegia: a model for the study of immobilization osteoporosis. *J. Bone Miner. Res.* 1988; 3: 305–10.
- Crabtree N, Reeve J. et al. Ambulatory level and asymmetrical weight bearing after stroke affects bone loss in the upper and lower part of the femoral neck differently: bone adaptation after decreased mechanical loading. *Bone*. 2000; 27: 701–7.
- Hamdy R. et al. Changes in bone mineral content and density after stroke. *Am. J. Phys. Med. Rehabil.* 1993; 72: 188–91.
- Pappone N, Mandes M. et al. Determinants of bone mineral density in immobilization: a study on hemiplegic patients. *Osteoporos. Int.* 1996; 6: 50–4.
- Sato Y, Maruoka H. et al. Development of osteopenia in the hemiplegic finger in patients with stroke. *Eur. Neurol.* 1996; 36: 278–83.
- Compston J, Cooper C. Bone densitometry in clinical practice. *BMJ*. 1995; 310: 1507–10.
- Ensrud K. et al. Hip and calcaneal bone loss increase with advancing age: longitudinal results from the study of osteoporotic fractures. *J. Bone Miner. Res.* 1995; 10: 1778–87.
- Jones G, Nguyen T. et al. Progressive loss of bone in the femoral neck in elderly people: longitudinal findings from the Dubbo osteoporosis epidemiology study. *BMJ*. 1994; 309: 691–5.
- Brown S, Rosen C. Osteoporosis. *Med. Clin. North. Am.* 2003; 87: 1039–63.
- Sato Y, Fujimatsu Y, Kikuyama M. et al. Influence of immobilization on bone mass and bone metabolism in hemiplegic elderly patients with a long-standing stroke. *J. Neurol. Sci.* 1998; 156: 205–10.
- Sato Y. Abnormal bone and calcium metabolism in patients after stroke. *Arch. Phys. Med. Rehabil.* 2000; 81:117–21.
- Sato Y, Kuno H, Kaji M. et al. Increased bone resorption during the first year after stroke. *Stroke*. 1998; 29:1373–7.
- Chantraine A, Nusgens B, Lapierre C. Bone remodeling during the development of osteoporosis in paraplegia. *Calcif. Tis. Int.* 1986; 38:323–7.
- Sato Y, Maruoka H, Oizumi K. et al. Vitamin D deficiency and osteopenia in the hemiplegic limbs of stroke patients. *Stroke*. 1996; 27:2183–7.