The Moldovan Medical Journal, February 2017, Vol. 60, No 1

DOI: 10.5281/zenodo.1050325
UDC: 618.39-021.3:577.112.386

OPEN ACCESS

Homocysteine and recurrent miscarriage

Visternicean Elena

Department of Obstetrics and Gynecology No 2
Nicolae Testemitsanu State University of Medicine and Pharmacy, Chisinau, the Republic of Moldova
Corresponding author: mecineanuelena@yahoo.com. Received December 26, 2016; accepted February 06, 2017

References

1. Bergen N. et al. Homocysteine and folate concentrations in early pregnancy and the risk of adverse pregnancy outcomes: the Generation R Study. BJOG: an international journal of obstetrics and gynaecology, 2012, 119 (6), 739 – 751.
2. Cao Y. et al. The association of idiopathic recurrent early pregnancy loss with polymorphisms in folic acid metabolism-related genes. Genes & Nutrition, 2014, 9: 402, 1-8.
3. Di Simone. et al. Effect of folic acid on homocysteine-induced trophoblast apoptosis. In: Molecular Human Reproduction, 2004, 10 (9), 665 – 669.
4. Furness D. et al. Folate,Vitamin B12,Vitamin B6 and homocysteine: impact on pregnancy outcome. Maternal and Child Nutrition, 2011, 1-12.
5. Gris J. et al. Antiphospholipid/antiprotein antibodies, hemostasis-related autoantibodies, and plasma homocysteine as risk factors for a first early pregnancy loss: a matched case-control study. Blood, 2003, 10, 3502 – 3513.
6. Kumar K. et al. Plasma homocysteine levels correlated to interactions between folate status and methylene tetrahydrofolate reductase gene mutation in women with unexplained recurrent pregnancy loss. Journal of obstetrics and gynaecology: the journal of the Institute of Obstetrics and Gynaecology, 2003, 23, 55 – 58.
7. Luciene de Souza Venancio, Roberto Carlos Burini, Winston Bonetti Yoshida. Dietary treatment of hyperhomocysteinemia in peripheral arterial disease. Jornal Vascular Brasileiro, 2010, 9 (1), 28-41.
8. Nelen W. et al. Homocysteine and folate levels as risk factors for recurrent early pregnancy loss. Obstetrics and Gynecology, 2000, 95, 519 – 524.
9. OpenEpi 2.3.1 – Open Source Epidemiologic Statistics for Public Health, http://www.openepi.com (visited 16.11.2016)
10. Refsum H. et al. Facts and Recommendations about Total Homocysteine Determinations: An Expert Opinion. Clinical Chemistry, 2004, 50 (1), 3- 32.
11. Refsum Helga. Folate, vitamin B12 and homocysteine in relation to birth defects and pregnancy outcome. British Journal of Nutrition, 2001, 85, Suppl. 2, S109-S113.
12. Ronnenberg A. et al. Preconception homocysteine and B vitamin status and birth outcomes in Chinese women. American Journal of Clinical Nutrition, 2002, 76, 1385 – 1391.
13. Stanger O. et al. Clinical use and rational management of homocysteine, folic acid, and B vitamins in cardiovascular and thrombotic diseases. In: Zeitschrift für Kardiologie, 2004, 93, 439 – 453.
14. Ueland P., Vollset S. Homocysteine and Folate in Pregnancy. Clinical Chemistry, 2004, 8, 1293 – 1295.
15. Urban J. et al. Serum homocysteine and nitric oxide levels in pregnancy complicated with intrauterine fetal growth restriction. Archives of Perinatal Medicine, 2007, 13 (3), 27 – 29.
16. Vollset S. et al. Plasma total homocysteine, pregnancy complications, and adverse pregnancy outcomes: the Hordaland Homocysteine Study. American Journal of Clinical Nutrition, 2000, 71, 962 – 968.
17. WHO. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva, World Health Organization, 2016. http://apps.who.int/iris/bitstream/10665/250796/1/9789241549912-eng.pdf?ua=1 (visited 20.12.2016).